O.D.260 Inc.

Product Information

pAd1127-11

Research use only

Catalog No: QP-29
Lot No: 1002

Contents:
pAd1127-11 plasmid DNA, 20 μg,
1 μg/μL in TE pH 7.5

Storage: -20°C

Features and Applications:

pAd1127-11 is a plasmid designed for constructing adenovirus vectors expressing transgenes under the control of a CMV promoter located in place of the E1 region of the Ad5 genome. It is a derivative of pAd1127-06, in which a cassette containing a CMV promoter- MCS- bovine growth hormone (bGH) polyA signal was inserted between the *Xba*I and *Acc*65I sites in clockwise orientation, i.e. towards the right end of the adenovirus genome. It contains *Pac*I and *Swa*I sites flanking the first 440 base pairs from the Ad5 genome (including the left ITR and packaging signal), the CMV-bGHpA cassette, and the pIX coding region. The sequences encompassing the kanamycin-resistance gene, the λ cos site, the adenovirus 0-1.3 map units, the CMV expression cassette and the pIX coding sequence are flanked by two SfiI restriction sites. These sites generate non-symmetrical sticky ends suitable for directional cloning with the other AdenoQuick2.0 plasmids (pAd1128, pAd1129, pAd1130, and their derivatives).
The packaging signal of pAd1127-11 extends to psn 440 (in the Ad5 genome), to include all 7 packaging "A" repeats (I, II, III, IV, V, VI, and VII). The complete packaging signal region might confer a growth advantage to the virus, according to Youil et al (Human Gene Therapy 14: 1017-1034). Because of the size of the E1 deletion (440-3510), the vectors generated from pAd1127-06 have minimal or no homology with the Ad5 sequences inserted in the chromosome of the helper cells such as PER-C6, thereby minimizing the probability of RCA generation. pAd1127-11 can also be used to manipulate the pIX promoter and coding region.

Selection:
prokaryotic – kanamycin 25 μg/mL

Replication:
prokaryotic – pUC ori



O.D.260 Inc.
PO Box 534, Boise, ID 83701
Ph. (208)345-7369 - FAX (208)345-7569
E-mail info@od260.com
www.od260.com